arXiv:math/0104031v1 [math.AG] 3 Apr 2001

Chern classes for representations of reductive groups

Arnaud Beauville

Introduction

Let G be a complex connected reductive group, and let R(G) be its representation ring. As an abelian group R(G) is spanned by the finite-dimensional representations [V] of G, with the relations $[\mathrm{V}\oplus\mathrm{W}]=[\mathrm{V}]+[\mathrm{W}]$; the ring structure is defined by the tensor product. Moreover the exterior product of representations give rise to a sequence of operations $\lambda^p:\mathrm{R(G)}\rightarrow\mathrm{R(G)}$ which make R(G) into a *λ-ring* – see (1.1) below for the definition.

In the course on his work on the Riemann-Roch theorem, Grothendieck had the remarkable insight that this purely algebraic structure is enough to define Chern classes, without any reference to a cohomology or Chow ring. He associated to any λ-ring R a filtration of R, the *γ-filtration*; the Chern classes take value in the associated graded ring $\mathrm{gr}\,\mathrm{R}$, and the Chern character is a ring homomorphism $\mathrm{ch}:\mathrm{R}\rightarrow\prod_p \mathrm{gr}^p_{\mathbf{Q}}$, where $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{R}=\mathrm{gr}\,\mathrm{R}\otimes\mathbf{Q}$. When applied to the Grothendieck ring K(X) of vector bundles on a smooth algebraic variety X these definitions give back the classical ones, at least modulo torsion: the graded ring $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{K(X)}$ coincides with the Chow ring $\mathrm{CH(X)}_{\mathbf{Q}}$, and the Chern classes with the usual ones.

The aim of this note is to compute the Chern classes for the representation ring R(G) – a simple exercise which I have been unable to find in the literature. Let $\mathfrak{g}$ be the Lie algebra of G; we will assume that G, and therefore $\mathfrak{g}$, are defined over $\mathbf{Q}$ (alternatively, we could without any loss take our Chern classes in $\mathrm{gr}_{\mathbf{C}}\,\mathrm{R(G)}$). We denote by $\mathrm{Pol}(\mathfrak{g})^{\mathrm{inv}}$ the ring of polynomial functions on $\mathfrak{g}$ which are invariant under the action of the adjoint group.

Theorem.– a) *The graded ring* $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{R(G)}$ *is canonically isomorphic to the ring* $\mathrm{Pol}(\mathfrak{g})^{\mathrm{inv}}$ *of invariant polynomial functions on* $\mathfrak{g}$.

b) *Let* ρ *be a representation of* G, *and* $\mathrm{L}\rho$ *the corresponding representation of* $\mathfrak{g}$. *The total Chern class* $c(\rho)$ *is equal to the invariant function* $\det(1+\mathrm{L}\rho)$, *and the Chern character* $\mathrm{ch}(\rho)$ *to* $\mathrm{Tr}(\exp\mathrm{L}\rho)$.

We have of course an explicit description of the ring R(G). Let T be a maximal torus of G; the Weyl group W acts on the ring R(T), and the restriction map $\mathrm{R(G)}\rightarrow\mathrm{R(T)}$ identifies R(G) with the invariant sub-ring $\mathrm{R(T)}^{\mathrm{W}}$. As an abelian group $\mathrm{R(T)}$ is spanned by one-dimensional elements (we will say that it is a *split* λ-ring), which makes easy to compute its γ-filtration and Chern classes. The theorem follows easily once we know that the γ-filtration of $\mathrm{R(T)}$ induces that of $\mathrm{R(G)}$. This turns out to be a general fact for invariant sub-rings of split λ-rings; we will deduce it from the behaviour of the γ-filtration under the Adams operations (Proposition 1.6).

In section 3 we discuss an application. Let P be a principal G-bundle over a base B (which may be a variety, or an arbitrary topos); the Theorem provides a simple definition of the characteristic classes of P in the graded ring $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{K(B)}$, and a simple way of computing the Chern classes of the associated vector bundles.

1. Generalities on λ-rings

In this section we recall the definition and basic properties of λ-rings. Standard references are [SGA6] or [F-L]; we follow the terminology of [SGA6], Exposé V.

(1.1) A λ-*ring* R is a commutative ring with two more pieces of structure:

– An augmentation, that is a ring homomorphism $\varepsilon : \mathrm{R} \to \mathbf{Z}$;

– A λ-structure, that is a sequence of maps $\lambda^i : \mathrm{R} \to \mathrm{R}$ such that, for any x, y in R and $n \in \mathbf{N}$,

$$\lambda^0(x) = 1 \qquad \lambda^1(x) = x \qquad \lambda^n(x+y) = \sum_{p+q=n} \lambda^p(x)\,\lambda^q(y)\ .$$

If we put $\lambda_t(x) = \sum_p \lambda^p(x)\,t^p \in \mathrm{R}[[t]]$, the last condition is equivalent to

$$\lambda_t(x+y) = \lambda_t(x)\,\lambda_t(y)\ .$$

Moreover we want formulas giving $\lambda^p(xy)$ and $\lambda^p(\lambda^q(x))$ as polynomials in $\lambda^1(x), \ldots, \lambda^p(x); \lambda^1(y), \ldots, \lambda^p(y)$ and $\lambda^1(x), \ldots, \lambda^{pq}(x)$ respectively. A convenient way of expressing these is to introduce a sequence $(\psi^k)_{k\geq 1}$ of additive endomorphisms of R, the *Adams operations*, defined by

$$\sum_{k\geq 1} \psi^k(x)\,(-t)^{k-1} = \lambda_t(x)^{-1}\,\frac{d}{dt}\lambda_t(x)\ .$$

Then the condition on the λ-structure means that the ψ^k are ring endomorphisms, and satisfy $\psi^k \circ \psi^\ell = \psi^{k\ell}$ for $k, \ell \geq 1$.

(1.2) Grothendieck associates to this situation a second λ-structure, defined by $\gamma_t(x) = \lambda_{\frac{t}{1-t}}(x)$, and a decreasing filtration of R, the γ-*fitration* $(\Gamma^p(\mathrm{R}))_{p\geq 0}$: we put $\Gamma^0 = \mathrm{R}$, $\Gamma^1 = \operatorname{Ker} \varepsilon$, and Γ^p is spanned by the elements $\gamma^{i_1}(x_1) \ldots \gamma^{i_k}(x_k)$

with $x_1, \dots, x_k \in \Gamma^1$ and $i_1 + \cdots + i_k = p$. Let $\mathrm{gr}\,\mathrm{R} = \bigoplus_{p\geq 0} \Gamma^p/\Gamma^{p+1}$ be the associated graded ring. The *Chern classes* $(c_p(x))_{p\geq 0}$ of an element $x \in \mathrm{R}$ are defined by

$$c_p(x) = \gamma^p(x - \varepsilon(x)) \quad \text{in} \quad \Gamma^p/\Gamma^{p+1} \ .$$

(1.3) A crucial point in what follows will be the behaviour of the γ-filtration with respect to the Adams operations. Let us say that an element x of R has λ-*dimension* n if it satisfies $\varepsilon(x) = n$ and $\lambda^i(x) = 0$ for $i > n$; we say that R is $\mathbf{Q}$-*finite* λ-*dimensional* if the $\mathbf{Q}$-vector space $\mathrm{R} \otimes \mathbf{Q}$ is spanned by elements of finite λ-dimension. For such a λ-ring we have, for each $k \geq 1$ and $x \in \Gamma^p(\mathrm{R})$:

$$\psi^k(x) \equiv k^p x \quad \text{mod. } \mathbf{Q}\Gamma^{p+1}(\mathrm{R}) \tag{1.4}$$

([F-L], III, Proposition 3.1).

(1.5) Let us say that the λ-ring R is *split* if is generated as an abelian group by elements of λ-dimension 1. In this case we have $\Gamma^p(\mathrm{R}) = \mathfrak{r}^p$, where $\mathfrak{r}$ is the augmentation ideal of R: indeed we have $\mathfrak{r}^p \subset \Gamma^p$ because γ^1 is the identity, and $\gamma_t(x) \in \sum_{p\geq 0} \mathfrak{r}^p\, t^p$ for all $x \in \mathfrak{r}$ because this holds for $x + y$ if it holds for x and y, and $\gamma_t(\xi - 1) = 1 + (\xi - 1)t$ if ξ has λ-dimension 1.

Proposition 1.6 .– *Let* R *be a split* λ*-ring, and* G *a finite group of automorphisms of the* λ*-ring* R. *Then the* G*-invariant elements form a sub-*λ*-ring* S *of* R. *The filtrations* $(\Gamma^p(\mathrm{S}))_{p\geq 0}$ *and* $(\Gamma^p(\mathrm{R}) \cap \mathrm{S})_{p\geq 0}$ *coincide in* $\mathrm{S} \otimes \mathbf{Q}$.

Proof: Since the action of G commutes with the λ^i the λ-structure of R induces a λ-structure on S. Moreover the $\mathbf{Q}$-vector space $\mathrm{S} \otimes \mathbf{Q}$ is spanned by the elements $\sum_{g\in \mathrm{G}} g\xi$, where ξ is an element of R of λ-dimension 1; these elements have λ-dimension $|\mathrm{G}|$, thus S is $\mathbf{Q}$-finite λ-dimensional.

To alleviate the notation we write R, S instead of $\mathrm{R} \otimes \mathbf{Q}$ and $\mathrm{S} \otimes \mathbf{Q}$, and $\Gamma^p(\mathrm{R}), \Gamma^p(\mathrm{S})$ instead of $\mathbf{Q}\Gamma^p(\mathrm{R}), \mathbf{Q}\Gamma^p(\mathrm{S})$. We have $\Gamma^p(\mathrm{S}) \subset \Gamma^p(\mathrm{R}) \cap \mathrm{S}$ for each p. Let us first prove that the two filtrations define the same topology. Let $\mathfrak{r} = \Gamma^1(\mathrm{R})$ be the augmentation ideal of R, and $\mathfrak{s} = \mathrm{S} \cap \mathfrak{r}$. Since R is a finite S-module, the ring $\mathrm{R}/\mathfrak{s}\mathrm{R}$ is artinian; thus there exists an integer ν such that $\mathfrak{r}^\nu \subset \mathfrak{s}\mathrm{R}$, and therefore $\mathfrak{r}^{n\nu} \subset \mathfrak{s}^n\mathrm{R}$ for all n.

As S-modules S is a direct summand of R (consider the projector $r \mapsto \dfrac{1}{|\mathrm{G}|}\sum_{g\in \mathrm{G}} gr$). Thus for every ideal $\mathfrak{a}$ of S, the induced homomorphism $\mathrm{S}/\mathfrak{a} \to \mathrm{R}/\mathfrak{a}\mathrm{R}$ is injective, which means $\mathfrak{a}\mathrm{R} \cap \mathrm{S} = \mathfrak{a}$. Using (1.5) we obtain

$$\Gamma^{p\nu}(\mathrm{R}) \cap \mathrm{S} = \mathfrak{r}^{p\nu} \cap \mathrm{S} \subset \mathfrak{s}^p\mathrm{R} \cap \mathrm{S} = \mathfrak{s}^p \subset \Gamma^p(\mathrm{S}) \ .$$

Let us prove now the inclusion $\Gamma^p(\mathrm{R}) \cap \mathrm{S} \subset \Gamma^p(\mathrm{S})$ by induction on p, the case $p = 0$ being obvious. We have just seen that there exists an integer N such that $\Gamma^{\mathrm{N}}(\mathrm{R}) \cap \mathrm{S} \subset \Gamma^p(\mathrm{S})$; let n be the smallest integer with that property. Assume $n > p$. Let $x \in \Gamma^{n-1}(\mathrm{R}) \cap \mathrm{S}$, and let $k \geq 2$ be an integer; by (1.4) we have

$$\psi^k(x) \equiv k^{n-1}x \quad (\text{mod. } \Gamma^n(\mathrm{R}) \cap \mathrm{S}) \ .$$

On the other hand we have $x \in \Gamma^{p-1}(\mathrm{S})$ by the induction hypothesis; since S is $\mathbf{Q}$-finite λ-dimensional (1.4) gives

$$\psi^k(x) \equiv k^{p-1}x \quad (\text{mod. } \Gamma^p(\mathrm{S})) \ .$$

Since $\Gamma^n(\mathrm{R}) \cap \mathrm{S} \subset \Gamma^p(\mathrm{S})$ we get $x \in \Gamma^p(\mathrm{S})$ for all $x \in \Gamma^{n-1}(\mathrm{R}) \cap \mathrm{S}$, contradicting the choice of n. Therefore we have $n \leq p$, hence $\Gamma^p(\mathrm{R}) \cap \mathrm{S} = \Gamma^p(\mathrm{S})$. ■

2. The $\boldsymbol{\lambda}$-ring R(G)

(2.1) Let T be a connected multiplicative group, and X its character group; this is a free finitely generated abelian group. The representation ring $\mathrm{R(T)}$ is isomorphic to the group algebra $\mathbf{Z}[\mathsf{X}]$; we denote by $([\alpha])_{\alpha \in \mathsf{X}}$ its canonical basis. The augmentation and the λ-structure are characterized by $\varepsilon([\alpha]) = 1$ and $\lambda_t([\alpha]) = 1 + t\alpha$.

Let $\mathfrak{r}$ be the augmentation ideal in $\mathrm{R(T)}$. We have a canonical isomorphism $\mathsf{X} \to \mathfrak{r}/\mathfrak{r}^2$ which maps a character α to the class of $[\alpha] - 1$ (observe that $[\alpha\beta] - 1 \equiv ([\alpha] - 1) + ([\beta] - 1)$ (mod. $\mathfrak{r}^2$). Since the rings $\mathrm{R(T)}$ and $\mathrm{R(T)}/\mathfrak{r} \cong \mathbf{Z}$ are regular, the canonical map $\mathsf{S}(\mathfrak{r}/\mathfrak{r}^2) \to \underset{p \geq 0}{\oplus} \mathfrak{r}^p/\mathfrak{r}^{p+1}$ is bijective; using (1.5) we get a canonical isomorphism

$$\varphi : \mathsf{S}(\mathsf{X}) \xrightarrow{\sim} \mathrm{gr}\, \mathrm{R(T)} \ .$$

Under this isomorphism the element $\gamma^1([\alpha] - 1)$ of $\mathfrak{r}/\mathfrak{r}^2$ corresponds to $\alpha \in \mathsf{X} \subset \mathsf{S}(\mathsf{X})$. Thus the total Chern class $c(\rho)$ of an element $\rho = [\alpha_1] + \cdots + [\alpha_d]$ is given by $c(\rho) = \prod_i (1 + \alpha_i)$.

Let $\mathfrak{h}$ be the Lie algebra of T, viewed as a vector space over $\mathbf{Q}$. We have a canonical isomorphism $\mathsf{X} \otimes \mathbf{Q} \to \mathfrak{h}^*$, which associates to a character $\alpha : \mathrm{T} \to \mathbf{G}_m$ its derivative $\mathrm{L}\alpha : \mathfrak{h} \to \mathbf{Q}$. Thus we can identify $\mathsf{S}(\mathsf{X}) \otimes \mathbf{Q}$ to the algebra $\mathrm{Pol}(\mathfrak{h})$ of polynomial maps $\mathfrak{h} \to \mathbf{Q}$; the above formula becomes $c(\rho) = \prod_i (1 + \mathrm{L}\alpha_i)$ in $\mathrm{Pol}(\mathfrak{h})$.

For $\alpha \in \mathsf{X}$, let $\mathbf{C}_\alpha$ denote the one-dimensional representation with character α. The element $\rho = [\alpha_1] + \ldots + [\alpha_d]$ of $\mathrm{R(T)}$ is the class of the representation $\mathrm{V} = \mathbf{C}_{\alpha_1} \oplus \ldots \oplus \mathbf{C}_{\alpha_d}$. In the corresponding representation $\mathrm{L}\rho$ of $\mathfrak{h}$, an element H

of $\mathfrak{h}$ acts through the diagonal matrix $\operatorname{diag}(\mathrm{L}\alpha_1(\mathrm{H}),\ldots,\mathrm{L}\alpha_d(\mathrm{H}))$. Thus $c(\rho)$ is equal to the function $\det(1+\mathrm{L}\rho)$ in $\mathrm{Pol}(\mathfrak{h})$.

The Chern character gives an homomorphism ch of $\mathrm{R(T)}$ into the ring $\widehat{\mathrm{Pol}}(\mathfrak{h})$ of formal series on $\mathfrak{h}$. We have $\mathrm{ch}([\alpha]) = e^{c_1([\alpha])} = e^{\mathrm{L}\alpha}$, hence

$$\mathrm{ch}(\rho) = e^{\mathrm{L}\alpha_1} + \ldots + e^{\mathrm{L}\alpha_d} = \mathrm{Tr}(\exp \mathrm{L}\rho) \quad \text{in } \widehat{\mathrm{Pol}}(\mathfrak{h}) \ .$$

Remark 2.2 .– The exponential morphism of formal groups $\exp : \widehat{\mathfrak{h}} \to \widehat{\mathrm{T}}$ induces an injective homomorphism $\exp^* : \mathcal{O}_{\widehat{\mathrm{T}},1} = \mathbf{Q}[[\mathsf{X}]] \to \mathcal{O}_{\widehat{\mathfrak{h}},0} = \widehat{\mathrm{Pol}}(\mathfrak{h})$, which maps $[\alpha]$ to $e^{\mathrm{L}\alpha}$. The Chern character is the composition of this map with the injection $\mathrm{R(T)} = \mathbf{Z}[\mathsf{X}] \hookrightarrow \mathbf{Q}[[\mathsf{X}]]$.

(2.3) Let G be a complex connected reductive group, $\mathfrak{g}$ its Lie algebra, T a maximal torus of G and $\mathfrak{h}$ its Lie algebra; we can assume that G , T , $\mathfrak{g}$ and $\mathfrak{h}$ are defined over $\mathbf{Q}$. The Weyl group W acts on T , hence on the ring R(T) ; restriction to T induces a homomorphism of λ-rings $\mathrm{R(G)} \to \mathrm{R(T)}$, whose image is the invariant sub-ring $\mathrm{R(T)}^{\mathrm{W}}$ ([SGA6], Exposé 0 App., Th. 1.1). By Proposition 1.6, the graded ring $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{R(G)}$ is isomorphic to $(\mathrm{gr}_{\mathbf{Q}}\,\mathrm{R(T)})^{\mathrm{W}}$, that is to the ring $\mathrm{Pol}(\mathfrak{h})^{\mathrm{W}}$ of invariant polynomials on $\mathfrak{h}$ (2.1).

Let ρ be a representation of G . From the commutative diagrams

$$\begin{tikzcd} \mathrm{R(T)} \arrow[r, "c"] & \widehat{\mathrm{Pol}}(\mathfrak{h}) \\ \mathrm{R(G)} \arrow[u] \arrow[r, "c"] & \widehat{\mathrm{Pol}}(\mathfrak{h})^{\mathrm{W}} \arrow[u] \end{tikzcd} \qquad \begin{tikzcd} \mathrm{R(T)} \arrow[r, "\mathrm{ch}"] & \widehat{\mathrm{Pol}}(\mathfrak{h}) \\ \mathrm{R(G)} \arrow[u] \arrow[r, "\mathrm{ch}"] & \widehat{\mathrm{Pol}}(\mathfrak{h})^{\mathrm{W}} \arrow[u] \end{tikzcd}$$

we obtain

$$c(\rho) = \det(1+\mathrm{L}\rho) \ \text{ in } \ \mathrm{Pol}(\mathfrak{h})^{\mathrm{W}} \ ; \quad \mathrm{ch}(\rho) = \mathrm{Tr}(\exp \mathrm{L}\rho) \ \text{ in } \ \widehat{\mathrm{Pol}}(\mathfrak{h})^{\mathrm{W}} \ .$$

The restriction map $\mathrm{Pol}(\mathfrak{g}) \to \mathrm{Pol}(\mathfrak{h})$ induces an isomorphism $\mathrm{Pol}(\mathfrak{g})^{inv} \to \mathrm{Pol}(\mathfrak{h})^{\mathrm{W}}$ ([B], § 8, Th. 1). Since the functions $\det(1+\mathrm{L}\rho)$ and $\mathrm{Tr}(\exp \mathrm{L}\rho)$ are invariant, the above equalities hold as well in $\mathrm{Pol}(\mathfrak{g})^{inv}$ and $\widehat{\mathrm{Pol}}(\mathfrak{g})^{inv}$ respectively. This proves the theorem stated in the introduction.

3. Application: Chern classes of associated bundles

(3.1) Let B be an algebraic variety, and P a principal G-bundle over B (for what follows B could be as well a scheme, or even a topos). To any representation

$\rho : \mathrm{G} \to \mathrm{GL(V)}$ is associated a vector bundle $\mathrm{P}^{\rho} = \mathrm{P} \times^{\mathrm{G}} \mathrm{V}$ on B ; we define in this way a homomorphism of λ-rings $b_{\mathrm{P}}^{*} : \mathrm{R(G)} \to \mathrm{K(B)}$ [1] . In view of the isomorphism described above, it induces a homomorphism of graded rings

$$c_{\mathrm{P}} : \mathrm{Pol}(\mathfrak{g})^{inv} \to \mathrm{gr}_{\mathbf{Q}}\mathrm{K(B)} \ ,$$

called the characteristic homomorphism.

Let $\ell = \dim \mathrm{T}$. Recall that there are homogeneous functions $\mathrm{I}_1, \ldots, \mathrm{I}_\ell$ in $\mathrm{Pol}(\mathfrak{g})^{inv}$ such that $\mathrm{Pol}(\mathfrak{g})^{inv} = \mathbf{Q}[\mathrm{I}_1, \ldots, \mathrm{I}_\ell]$ ([B], § 8, Théorème 1). The elements $c_{\mathrm{P}}^{(i)} := c_{\mathrm{P}}(\mathrm{I}_i)$ may be called the *characteristic classes* of P (but note that they depend on the choice of the generating sequence $(\mathrm{I}_1, \ldots, \mathrm{I}_\ell)$). If B is a smooth variety, the graded ring $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{K(B)}$ is canonically isomorphic to the rational Chow ring $\mathrm{CH}_{\mathbf{Q}}(\mathrm{B})$ ([SGA6], Exposé XIV, n° 4); our characteristic classes correspond under this isomorphism to those defined in [V] and [E-G].

Proposition 3.2 .– *Let* ρ *be a representation of* G *; write* $\det(1 + \mathrm{L}\rho) = \mathrm{F}(\mathrm{I}_1, \ldots, \mathrm{I}_\ell)$ *, where* F *is a polynomial in* ℓ *indeterminates. Let* P *be a principal* G*-bundle on* B *, with characteristic classes* $c_{\mathrm{P}}^{(1)}, \ldots, c_{\mathrm{P}}^{(\ell)}$ *in* $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{K(B)}$ *. Then the total Chern class in* $\mathrm{gr}_{\mathbf{Q}}\,\mathrm{K(B)}$ *of the associated bundle* P^{ρ} *is*

$$c(\mathrm{P}^{\rho}) = \mathrm{F}(c_{\mathrm{P}}^{(1)}, \ldots, c_{\mathrm{P}}^{(\ell)}) \ .$$

Similarly, if $\mathrm{Tr}(\exp \mathrm{L}\rho) = \mathrm{G}(\mathrm{I}_1, \ldots, \mathrm{I}_\ell)$ *, with* $\mathrm{G} \in \mathbf{Q}[[\mathrm{T}_1, \ldots, \mathrm{T}_\ell]]$ *, we have* $\mathrm{ch}(\mathrm{P}^{\rho}) = \mathrm{G}(c_{\mathrm{P}}^{(1)}, \ldots, c_{\mathrm{P}}^{(\ell)})$.

Proof : This follows from the Theorem, the commutative diagram

$$\begin{tikzcd}
\mathrm{R(G)} \arrow[r, "b_{\mathrm{P}}^{*}"] \arrow[d, "c"'] & \mathrm{K(B)} \arrow[d, "c"] \\
\widehat{\mathrm{Pol}}(\mathfrak{g})^{inv} \arrow[r, "c_{\mathrm{P}}"] & \widehat{\mathrm{gr}}_{\mathbf{Q}}\,\mathrm{K(B)}
\end{tikzcd}$$

and the corresponding diagram for the Chern character. ■

Examples 3.3 .– The Proposition provides a way of computing the characteristic classes in terms of Chern classes, at least for classical groups. We use the standard generating system $(\mathrm{I}_1, \ldots, \mathrm{I}_\ell)$ given for instance in [B], § 13. We denote by E_{P} the vector bundle on B associated to P through the standard representation of $\mathrm{GL}(n)$, $\mathrm{SO}(n)$ or $\mathrm{Sp}(n)$ in $\mathbf{C}^n$.

---

[1] In fancy terms, the principal bundle P corresponds to a morphism b_{P} of B into the classifying topos (or stack) BG , and b_{P}^{*} is just the pull-back map.

a) For $G = GL(\ell)$, we have $I_p(A) = \mathrm{Tr}(\mathsf{\Lambda}^p A)$ for $A \in \mathfrak{gl}(\ell)$ and $1 \leq p \leq \ell$; this gives $c_P^{(p)} = c_p(E_P)$ for $1 \leq p \leq \ell$.

b) For $G = \mathrm{Sp}(2\ell)$ or $\mathrm{SO}(2\ell+1)$, we have $I_p(A) = \mathrm{Tr}(\mathsf{\Lambda}^{2p} A)$; this gives $c_P^{(p)} = c_{2p}(E_P)$ for $1 \leq p \leq \ell$ (the vector bundle E_P is isomorphic to its dual, thus its odd Chern classes with rational coefficients vanish).

c) For $G = \mathrm{SO}(2\ell)$, we realize $\mathfrak{g}$ as the space of skew-symmetric matrices in $\mathfrak{gl}(2\ell)$; we have $I_p(A) = \mathrm{Tr}(\mathsf{\Lambda}^{2p} A)$ for $p < \ell$, and $I_{2\ell}(A) = \mathrm{Pf}(A)$. We obtain $c_P^{(p)} = c_{2p}(E_P)$ for $1 \leq p \leq \ell - 1$, and $c_P^{(\ell)}$ is a class in $\mathrm{gr}^\ell_{\mathbf{Q}} K(B)$ with square $c_{2\ell}(E_P)$.

REFERENCES

[B] N. Bourbaki: *Groupes et algèbres de Lie*, chapitre VIII. Hermann, Paris (1975).

[E-G] D. Edidin, W. Graham: *Characteristic classes in the Chow ring*. J. Algebraic Geom. **6** (1997), 431–443.

[F-L] W. Fulton, S. Lang: *Riemann-Roch algebra*. Grundlehren der Math. **277**. Springer-Verlag, New York (1985).

[SGA6] *Théorie des intersections et théorème de Riemann-Roch*. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6). Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Lecture Notes in Math. **225**, Springer-Verlag, Berlin-New York (1971).

[V] A. Vistoli: *Characteristic classes of principal bundles in algebraic intersection theory*. Duke Math. J. **58** (1989), 299–315.

Arnaud Beauville
Laboratoire J.-A. Dieudonné
UMR 6621 du CNRS
Université de Nice
Parc Valrose
F-06108 Nice Cedex 2